

Research Review
NEW EVIDENCE ON FLUORIDATION

M Diesendorf, J Colquhoun, B J Spittle, D N Everingham and F W Clutterbuck
Sydney and Brisbane, Australia; Auckland and Dunedin, New Zealand

Reprinted, with permission, from *The Australian and New Zealand
Journal of Public Health* Vol. 21 No. 2 1997

ABSTRACT: A review of recent scientific literature reveals a consistent pattern of evidence – hip fractures, skeletal fluorosis, the effect of fluoride on bone structure, fluoride levels in bones and osteosarcomas – pointing to the existence of causal mechanisms by which fluoride damages bones. In addition, there is evidence, accepted by some eminent dental researchers and at least one leading US proponent of fluoridation, that there is negligible benefit from ingesting fluoride, and that any (small) benefit from fluoridation comes from the action of fluoride at the surface of the teeth before fluoridated water is swallowed. Public health authorities in Australia and New Zealand have appeared reluctant to consider openly and frankly the implications of this and earlier scientific evidence unfavourable to the continuation of the fluoridation of drinking water supplies.

In recent years, new scientific evidence has emerged which suggests that there are significant risks and negligible benefits from ingesting low levels of fluoride. We outline the evidence that fluoridation of water supplies is harmful to bone, while providing negligible benefits when swallowed.

In focussing on the new evidence (mostly since 1989) in just two areas, it is not intended to diminish the importance of earlier evidence for concern about the health hazards of fluoridation: notably dental fluorosis, allergies and intolerance reactions, and genetic damage. These are reviewed elsewhere.¹⁻³

Mark Diesendorf is Professor of Environmental Science and Director, Institute for Sustainable Futures, University of Technology, Sydney, PO Box 123, Broadway NSW 2007, Australia. Fax 61 2 9209 4351. John Colquhoun is an Honorary Research Fellow, School of Education, University of Auckland. Bruce Spittle is Senior Lecturer, Department of Psychological Medicine, University of Otago School of Medical, Dunedin. Douglas Everingham is a retired medical practitioner and former Australian Federal Minister of Health. Frederick Clutterbuck is a medical practitioner in Brisbane.

Correspondence to Professor Diesendorf.

Fluoride damages bones

Since 1990, five major epidemiological studies from three countries – the United States (US), United Kingdom and France – showing a higher rate of hip fractures in fluoridated regions than unfluoridated regions have been reported in leading peer-reviewed journals.⁴⁻⁸ Although two of these reports were published as letters, the first was a correction to a refereed publication⁹ and the second was a supplement to a refereed publication about a prospective study which took account of major individual risk factors.¹⁰ In addition, a prospective study from the US shows a higher rate of hip fractures in a region naturally fluoridated with four parts per million (ppm) fluoride in drinking water than in a comparison region with 1 ppm.¹¹ Although there have been a few studies that have found no difference between fluoridated and unfluoridated regions, they have been either limited to small samples, or the women were not exposed to fluoride during the time of their lives when fluoride would be expected to affect bone most, that is, before menopause.¹²

The main weight of the recent evidence on hip fractures is consistent with earlier evidence from naturally fluoridated areas that low levels of fluoride ingested for several decades can cause the disease of bones and joints known as osteofluorosis or skeletal fluorosis. Evidence of skeletal fluorosis has been reported in at least nine studies from five countries with fluoride concentrations in drinking water of 0.7 to 2.5 ppm. These studies, and the inadequacies of studies that assert that there is no skeletal fluorosis in the US at fluoride concentrations below 4 ppm, have been reviewed elsewhere.²

In three to four decades, when people in areas where water is artificially fluoridated have accumulated fluoride in their bones from birth to old age, the increase in rates of hip fractures and skeletal fluorosis will be larger.

Fluoride has been used in high doses (20 to 32 mg a day) for short periods (one to two years) to treat osteoporosis. It is now recognised widely that, while this therapy adds mass to bones, it also damages the bone structure and leads to a higher risk of hip fracture.¹³⁻¹⁸ Bone analyses have shown that elderly women who lived for at least a decade in the town of Kuopio, Finland, with 1 ppm fluoride in its water supply, had high levels of fluoride in bone (typically 900 to 2300 ppm, but for women with impaired kidney function, as high as 3890 ppm).^{19,20} These levels are as high as have been reported in patients who have undergone fluoride therapy for osteoporosis.²¹

In the US National Cancer Institute's Surveillance, Epidemiology and End Results Program, an increase of 79 per cent was found in the incidence of osteosarcomas in young men living in fluoridated areas of Iowa and Seattle, but not in the unfluoridated areas, where the incidence decreased by 4 per cent.²² In fluoridated regions of the State of New Jersey, the incidence of osteosarcoma was three to seven times higher among males aged 10 to 19 years than in unfluoridated regions.²³ Osteosarcoma is a rare disease and so more evidence is required before any conclusions are drawn. But there is already a strong basis for concern, because the human data are supported by an animal experiment: the US National Toxicology Program has recorded a statistically significant, dose-related increase in the incidence of osteosarcoma in male rats ingesting fluoride.^{24,25}

Thus, there is a consistent pattern of evidence -- hip fractures, skeletal fluorosis, the effect of fluoride on bone structure, fluoride levels in bones, and osteosarcomas -- pointing to the existence of causal mechanisms by which fluoride damages bones.

Negligible benefit from fluoride ingestion

Recent research on the mechanism of action of fluoride in reducing the prevalence of dental caries (tooth decay) in humans shows that fluoride acts topically (at the surface of the teeth) and that there is negligible benefit in actually ingesting it.²⁶⁻²⁸ This is supported by experiments on laboratory rats: a slow-release source of fluoride fixed in the mouth reduced dental caries, but when the mouth was bypassed by placing the source under the skin, there was no detectable reduction.²⁹ The lack of observed systemic benefit from ingesting fluoridated water at a concentration 1 ppm is not surprising, since the resulting level of fluoride in the saliva is only around 0.01 ppm.³⁰

The evidence that there is negligible systemic benefit from fluoridation is accepted by eminent dental researchers²⁶⁻²⁸ and at least one leading US proponent of fluoridation, Professor Brian Burt.³¹ Therefore, proponents must come to grips with a serious ethical question: is it right to put fluoride in *drinking* water and to mislead the community that fluoride must be ingested, when any small benefit is due to the topical action of fluoride on teeth.³²

Alleged benefit from fluoride

We say 'any small benefit', because the results of recent large-scale studies in at least three countries show that, when similar communities are compared and the traditional *DMFT* (number of decayed, missing and filled teeth) index of dental caries is used, there is no detectable difference in caries prevalence. This has been demonstrated for schoolchildren in the major cities of New Zealand, Australia, the US and elsewhere.³³⁻³⁸ (When the newer *DMFS* (number of decayed, missing and filled surfaces) index was used, a 20% reduction was reported for US,³⁹ but, in absolute terms, this is only a fraction of a cavity per child.)

Of the many studies used by proponents of fluoridation to claim that there are enormous benefits from fluoridation, not one is a randomised controlled trial. Those that have been re-examined have been found to have serious design flaws.^{38,40-44} Indeed, hardly any of the many small-scale studies by enthusiasts of fluoridation are 'blind' and, in the rare cases when they are, the so-called 'control' was selected from a known high-caries area.⁴³ Many studies have also failed to take into account that unfluoridated towns tend to be rural, while fluoridated towns tend to be large cities, and that there is generally more dental caries in rural areas, irrespective of fluoridation status. In general, diet tends to be better in urban areas.

Many other studies have had no controls. Their authors have justified their profluoridation conclusions on the basis of large temporal declines in tooth decay. But, it is now known that equally large declines in caries have taken place in unfluoridated areas,⁴⁵⁻⁴⁸ and that in several cases this decline commenced before fluoride in any form was used to a significant degree.^{47,48}

However, there is now abundant evidence that topical uses of fluoride, extensively practised in Europe instead of water fluoridation, are effective in controlling tooth decay.⁴⁹ We agree that their cautious uses in dentistry are justified and provide an alternative to fluoridation which satisfies ethical concerns.³² However, in the past they have been promoted and practised rather irresponsibly – for example, the provision of highly concentrated fluoride toothpastes and mouth rinses to young children who inevitably ingest much of the fluoride. Too often overlooked is the evidence that tooth decay is associated with inadequate diets,⁵⁰ and that dietary control of caries, without the use of fluoride, is possible.⁵¹

Bias of health authorities

In our view, the evidence indicates that fluoridation entails real health risks and at best very small benefits. Therefore, the fluoridation of water supplies should be terminated forthwith. Yet, both in Australia and New Zealand, health authorities appear to be redoubling their efforts to fluoridate the remaining towns that have so far managed to hold fluoridation at bay.

The 1991 report on fluoridation by the National Health and Medical Research Council was published just as the first papers reporting the link between fluoridation and hip fractures were being published.⁵² It acknowledged in its section 6.4 some of the evidence that skeletal fluorosis is a potential health hazard, but created the false impression in its executive summary that there are no health risks. It is the executive summary which is read by decision-makers and the media. The report's profluoridation bias was further demonstrated by its failure to cite any of the studies presenting the evidence against fluoridation published in refereed journals.

The 1995 Report to the Minister by the New Zealand Public Health Commission demonstrated similar bias by failing even to cite any of the published papers on hip fractures, skeletal fluorosis or osteosarcomas.⁵³ However, the 1994 New Zealand Public Health Commission report did include some of these references and did acknowledge that:

It is possible that there is a small increased risk of hip fracture associated with water fluoridation, though the evidence for this is very inconclusive. More research is required to clarify this issue. A large amount of research has failed to provide evidence that exposure to fluoride causes cancer. However, the possibility of a small increased risk of osteosarcoma (a rare type of bone cancer) in young men cannot be ruled out at this stage. Here again, more research is needed. [From the executive summary; there are similar statements on p. 74 and p 78.]⁵⁴

But this information, and the references supporting it, were not forwarded officially to the Minister.

One of us (DE), while Federal Minister for Health in Australia from 1972 to 1975, could not get frank answers from his own department on the risks and benefits of fluoridation. Another of us (JC), while convenor of the New Zealand Fluoridation Promotion Committee, observed at first hand how his then fellow proponents of

fluoridation kept from the public and decision makers the evidence that fluoridation is less effective than claimed by proponents and is harmful, and then represented the evidence in a misleading way when it was eventually released. All of us have observed attempts by the medical and dental establishment in profluoridation countries to evade the evidence of concern and to suppress and misrepresent scientists, medical practitioners and dentists who attempt to publish evidence against fluoridation.⁵⁵⁻⁵⁷

For these and other reasons, we have no confidence in the impartiality of those institutions of government and the professions which have endorsed fluoridation for decades. Those who have built their careers and professional status on fluoridation cannot credibly assess the evidence against it. We have submitted this short paper for publication in the hope that at least some kind of scholarly debate will ensue.

REFERENCES

1. Waldbott GL, Burgstahler AW, McKinney HL. *Fluoridation: The Great Dilemma*. Coronado Press, Lawrence KS 1978.
2. Diesendorf M. The health hazards of fluoridation: a re-examination. *International Clinical Nutrition Review* 10 304-21 1990.
3. Yiamouyiannis J. *Fluoride: the Aging Factor* (3rd edition). Health Action Press, Delaware OH 1993.
4. Jacobsen SJ, Goldberg J, Miles TP *et al*. Regional variation in the incidence of hip fracture: US white women aged 65 years and older. *Journal of the American Medical Association* 264 500-502 1990.
5. Cooper C, Wickham CAC, Barker DJR, Jacobsen SJ. Water fluoridation and hip fracture [letter]. *Journal of the American Medical Association* 266 513-514 1991.
6. Danielson C, Lyon JL, Egger M, Goodenough GK. Hip fractures and fluoridation in Utah's elderly population. *Journal of the American Medical Association* 268 746-748 1992.
7. Jacobsen SJ, Goldberg J, Cooper C, Lockwood SA. The association between water fluoridation and hip fracture among white women and men aged 65 years and older: a national ecologic study. *Annals of Epidemiology* 2 617-626 1992.
8. Jacqmin-Gadda H, Commenges D, Dartigues J-F. Fluorine concentration in drinking water and fractures in the elderly [letter]. *Journal of the American Medical Association* 273 775-776 1995.
9. Cooper C, Wickham C, Lacey RF, Barker DJP. Water fluoride concentration and fracture of the proximal femur. *Journal of Epidemiology and Community Health* 44 17-19 1990.
10. Dartigues JF, Gagnon M, Barberger-Gateau P *et al*. The Paquid epidemiological program on brain aging. *Neuroepidemiology* 11 (Suppl 1) 14-18 1992.
11. Sowers MFR, Clark MK, Jannausch ML, Wallace RB. A prospective study of bone mineral content and fracture in communities with differential fluoride exposure. *American Journal of Epidemiology* 133 649-660 1991.
12. Lee JR. Fluoridation and hip fracture. According to the National Research Council Report: 'Health effects of ingested fluoride'. *Fluoride* 26 274-277 1993.

13. Hedlund LR, Gallagher JC. Increased incidence of hip fracture in osteoporotic women treated with sodium fluoride. *Journal of Bone and Mineral Research* 4 223-225 1989.
14. Lindsay R. Fluoride and bone – quantity versus quality (Editorial). *New England Journal of Medicine* 322 845-846 1990.
15. Melton LJ. Fluoride in the prevention of osteoporosis and fractures. *Journal of Bone and Mineral Research* 5 (Suppl 1) S163-167 1990.
16. Riggs BL, Hodgson SF, O'Fallon WM *et al.* Effect of fluoride treatment on the fracture rate in postmenopausal women with osteoporosis. *New England Journal of Medicine* 322 802-809 1990.
17. Fratzl P, Roschger P, Eschberger J *et al.* Abnormal bone mineralisation after fluoride treatment in osteoporosis: a small-angle x-ray-scattering study. *Journal of Bone and Mineral Research* 9 1541-1549 1994.
18. Sogaard CH, Mosekilde L, Richards A, Mosekilde L. Marked decrease in trabecular bone quality after five years of sodium fluoride therapy - assessed by biomechanical testing of iliac crest bone biopsies in osteoporotic patients. *Bone* 15 393-399 1994.
19. Arnala I. *Bone fluoride, histomorphometry and incidence of hip fracture.* University of Kuopio, Kuopio 1983.
20. Arnala I, Alhava EM, Kauranen P. Effects of fluoride on bone in Finland: histomorphometry of cadaver bone from low and high fluoride areas. *Acta Orthopædica Scandinavica* 56 161-166 1985.
21. Baud CA, Lagier R, Bang S *et al.* Treatment of osteoporosis with NaF, calcium or/and phosphorus, and vitamin D: histological, morphometric and biophysical study of the bone tissue. In: Courvoisier B, Donath A, Baud CA (Eds). *Fluoride and Bone.* Hans Huber, Berne 1978 pp 290-292.
22. Hoover RN, Devesa S, Cantor K, Fraumeni Jr JF. Time trends for bone and joint cancers and osteosarcomas in the Surveillance, Epidemiology and End Results (SEER) Program, National Cancer Institute. In: *Review of Fluoride: Benefits and Risks.* Report of the ad hoc committee on fluoride of the committee to coordinate environmental health and related programs. Department of Health and Human Services, Public Health Service, Washington DC 1991 pp F1-F7.
23. Cohn PD. *A brief report on the association of drinking water fluoridation and the incidence of osteosarcoma among young males.* New Jersey Department of Health, Trenton NJ 1992.
24. *The toxicology and carcinogenesis of sodium fluoride in F344/N rats and B6C3F1 mice* (National Toxicology Program technical report 393. Publication No. 90-2848). National Institutes of Health, Bethesda MD 1990.
25. Maurer JK, Cheng MC, Boysen BG, Anderson RL. Two-year carcinogenicity study of sodium fluoride in rats. *Journal of the National Cancer Institute* 82 1118-1126 1990.
26. Margolis HC, Moreno EC. Physicochemical perspectives on the cariostatic mechanisms of systemic and topical fluorides. *Journal of Dental Research* 69 (special issue) 606-613 1990.
27. Thylstrup A. Clinical evidence of the role of pre-eruptive fluoride in caries prevention. *Journal of Dental Research* 69 (special issue) 742-750 1990.
28. Comments by Fejerskov O, in Whitford GM (compiler). Discussion of Session I: Metabolism of fluoride. *Journal of Dental Research* 69 (special issue) 556-557 1990.

29. Mirth DB, Adderly DD, Monell-Torrens E *et al.* Comparison of the cariostatic effect of topically and systemically administered controlled-release fluoride in the rat. *Caries Research* 19 466-474 1985.
30. Oliveby A, Twetman S, Ekstrand J. Diurnal fluoride concentration in whole saliva in children living in a high- and a low-fluoride area. *Caries Research* 24 44-47 1990.
31. Debate between John R Lee and Brian A Burt. *Fluoride* 27 180-182 1994.
32. Diesendorf M. How science can illuminate ethical debates: a case study on water fluoridation. *Fluoride* 28 87-104 1995.
33. Colquhoun J. Child dental health differences in New Zealand. *Community Health Studies* 11 85-90 1987.
34. Gray AS. Fluoridation: Time for a new base line? *Journal of the Canadian Dental Association* 53 763-765 1987.
35. Colquhoun J. Is there a dental benefit from fluoridation? *Fluoride* 27 13-22 1994.
36. Hildebolt CF, Elvin-Lewis M, Molnar S *et al.* Caries prevalences among geochemical regions of Missouri. *American Journal of Physical Anthropology* 78 79-92 1989.
37. Yiamouyiannis JA. Water fluoridation and tooth decay: Results from the 1986-1987 national survey of US schoolchildren. *Fluoride* 23 55-67 1990.
38. Diesendorf M. Have the benefits of water fluoridation been overestimated? *International Clinical Nutrition Review* 10 292-303 1990.
39. Brunelle JA, Carlos JP. Recent trends in dental caries in US children and the effect of water fluoridation. *Journal of Dental Research* 69 (special issue) 723-727 1990.
40. Sutton PRN. *Fluoridation: Errors and Omissions in Experimental Trials* (2nd edition). Melbourne University Press, Melbourne 1960.
41. Diesendorf M. A re-examination of Australian fluoridation trials. *Search* 17 256-261 1986.
42. Colquhoun J, Mann R. The Hastings fluoridation experiment: Science or swindle? *The Ecologist* 16 (6) 243-248 1986, 17 (2) 125-126 1987.
43. Diesendorf M. Anglesey fluoridation trials re-examined. *Fluoride* 22 (2) 1989.
44. Colquhoun J. Flawed foundation: A re-examination of the scientific basis for a dental benefit from fluoridation. *Community Health Studies* 14 288-296 1990.
45. Glass RL (Ed). First international conference on the declining prevalence of dental caries. *Journal of Dental Research* 61 (special issue) 1304-1383 1982.
46. Leverett DH. Fluorides and the changing prevalence of dental caries. *Science* 217 26-30 1982.
47. Diesendorf M. The mystery of declining tooth decay. *Nature* 322 125-129 1986.
48. Colquhoun J. Decline in primary tooth decay in New Zealand. *Community Health Studies* 12 187-191 1988.
49. Fejerskov O, Ekstrand J, Burt BA (Eds). *Fluoride in Dentistry* (2nd edition) Munksgaard, Copenhagen 1996.
50. Price WA. *Nutrition and Physical Degeneration*. Hoeber, New York 1939.
51. Goldsworthy NE. Every doctor a dietitian. *Medical Journal of Australia* 1 285-286 1960.
52. National Health and Medical Research Council. *The Effectiveness of Water Fluoridation*. Australian Government Publishing Services, Canberra 1991.

53. *Fluoride and Oral Health: the Public Health Commission's advice to the Minister of Health*. New Zealand: Public Health Commission, Wellington 1995.
54. *Water Fluoridation In New Zealand: An Analysis and Monitoring Report*. New Zealand Public Health Commission, Wellington 1994.
55. Waldbott GL. *A Struggle With Titans*. Carlton Press, New York 1965
56. Varney W. *Fluoride in Australia: A Case to Answer*. Hale and Iremonger, Sydney 1986.
57. Martin B. *Scientific Knowledge in Controversy: The Social Dynamics of the Fluoridation Debate*. State University of New York Press, Albany 1991 (Chapter 5).

Reprinted, with permission, from *The Australian and New Zealand Journal of Public Health*
in *Fluoride Vol. 30* No. 3 1997, published by the International Society for Fluoride
Research